

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if # # adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ## child takes more than 5 doses in 24 hours ## taken with other drugs containing acetaminophen # adult has 3 or more alcoholic drinks every day while using this product

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have
before use if user has liver disease

Ask a doctor or pharmacist before use if you are
the user is taking the blood thinning drug warfarin

When using this product

Stop use and ask a doctor if
symptoms do not improve ## pain gets worse or lasts for more than 10 days # fever gets worse or lasts for more than 3 days # new symptoms occur ## redness or swelling is present # a rare sensitivity reaction occurs##
You may report side effects to 1-888-952-0050

If pregnant or breast-feeding

Keep out of reach of children
In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

Directions

Other information

Temporary relief of minor aches and pains associated with # common cold # headache # toothache # muscular aches # backache # arthritis # menstrual cramps # and reduction of fever
Adults and children 12 years of age and older: take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours;
Children 6-11 years of age: take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours;
Children under 6 years of age: do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.
store at room temperature 59°-86° F (15°-30° C) # do not use if imprinted safety seal under cap is broken or missing
corn starch, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolate.

Inactive ingredients